Breast Surgery

The following instructions are provided to assist you with planning your self care and recovery after your operation.

- 1. You must have a responsible adult accompany you home and stay with you overnight.
- 2. Plan on getting an adequate amount of rest for a few days following your operation. Return to normal activity gradually.
- 3. It is important to exercise the arm daily as per your doctor's instructions.
- 4. You are advised that you should not operate a motor vehicle, machinery, or perform any tasks requiring skill, co-ordination or judgment, or make any legal decisions within 24 hours after your operation.
- 5. Take all of your regular medications as you took them before your operation, unless instructed otherwise.
- 6. Take pain medications as prescribed. Try to avoid constipation by eating high fiber foods, drinking plenty of fluids and use a stool softener or mild laxative if needed. Your local pharmacist can be of assistance in choosing these products.
- 7. Resume your regular diet as tolerated. Do not drink any alcoholic beverages or use recreational drugs including beer and wine for 24 hours after your surgery.
- 8. It is important to have emptied your bladder (passed your urine) within 6 hours of returning home. If not, return to the Emergency Department of the nearest hospital.
- 9. You may notice that the solution used to cleanse your skin for surgery has coloured your skin. This will wash off with soap and water.
- 10. Keep fabric wrap on for 5 days. Once wrap is removed by Surgeon continue to wear a supportive bra for the next month including while sleeping.
- 11. If you have steri-strips (tapes) they will fall off over a 2 week period; please let them fall off on their own rather than removing them.
- 12. If you have a drain follow these instructions:
- 13. You should note the amount and character of the drainage when emptying the drain. You can then report this information to your doctor.



Breast Surgery

- 14. Do not allow injections, placement of IVs, drawing of blood or placement of a blood pressure cuff on the affected arm until advised by your doctor to do so.
- 15. You may shower after the dressing has been removed, gently pat incision dry.
- 16. Contact your Doctor or seek medical attention if any of the following occurs:
 - Signs of infection, including fever and chills
 - Redness, swelling, increasing pain, excessive bleeding, or discharge at the incision site.
 - Cough, shortness of breath, chest pain
 - Pain that you cannot control with the medication that you have been given.
 - Nausea and/or vomiting that you cannot control or which persist for more than two days after your surgery.
 - Pain and/or swelling in your feet, calves or legs.

If you have any concerns:

- ... Call your surgeon/doctor,
- ... or call Telehealth 1-866-797-0000
- ... or come to the Emergency Department at Georgian Bay General Hospital

17.The	lay after your surgery call your Surgeon's office to make week(s)	your follow-up appointment for
Specific	Instructions:	

